




PAIN-01-Peds: Percentage of pediatric patients < 18 years old who undergo a procedure and receive at least one non-opioid adjunct preoperatively or intraoperatively.



ECT
Endoscopy
Diagnostic Imaging Procedure
Obstetric Procedures
Procedure Type: Cardiac *Open Cardiac
TEE/Cardioversions
Block only cases
Non Operative Procedures (CPT: 00124, 00532, 00635, 01990, 01991, 01992, 01999)





Neuraxial Anesthesia: Caudal, Epidural and Neuraxial-Multiple determined by <u>Anesthesia Technique: Neuraxial</u>

<u>Analgesia Given</u>: Cases without any analgesia administered between Preop Start and Anesthesia End will be excluded.

Analgesia is defined as Peripheral Block, Neuraxial Block, Local Injection, Opioid Medication, or Opioid Sparing

Regional Block determined by <u>Anesthesia Technique</u>: <u>Peripheral Nerve Block</u>

OPIOID SPARING MEDICATION – must be administered between Anesthesia Start and Anesthesia End via a valid route.

10007 Acetaminophen 10222 Ibuprofen 10747 Naproxen

Medication

50380 'Emergence - ETT in place' documented between procedure end and anesthesia end

No

Sedation medication

Anesthesia End?

Patient had at least one non-opioid adjunct during the preoperative or intraop period?

No

Flag

Exclude

Exclude